



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2015

---

## **Lithium diisopropylamide (lda) as an efficient reducing agent for thioketones - mechanistic consideration**

Jasiński, Marcin ; Młostoń, Grzegorz ; Gebert, Andreas ; Heimgartner, Heinz

**Abstract:** Treatment of thiocarbonyl compounds with excess LDA leads to the corresponding thiols or sulfides depending on the work-up procedure. The mechanistic scenario for this unusual reduction pathway is discussed.

DOI: <https://doi.org/10.1080/10426507.2015.1012201>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-112069>

Journal Article

Accepted Version

Originally published at:

Jasiński, Marcin; Młostoń, Grzegorz; Gebert, Andreas; Heimgartner, Heinz (2015). Lithium diisopropylamide (lda) as an efficient reducing agent for thioketones - mechanistic consideration. *Phosphorus, Sulfur, and Silicon and the Related Elements*, 190:1281-1284.

DOI: <https://doi.org/10.1080/10426507.2015.1012201>


# LITHIUM DIISOPROPYLAMIDE (LDA) AS AN EFFICIENT REDUCING AGENT FOR THIOKETONES - MECHANISTIC CONSIDERATION

Marcin Jasiński,<sup>a</sup> Grzegorz Mlostoń,<sup>a\*</sup> Andreas Gebert,<sup>b</sup> and Heinz Heimgartner<sup>b</sup>

<sup>a</sup> Faculty of Chemistry, University of Łódź, Tamka 12, PL 91-403 Łódź, Poland

<sup>b</sup> Department of Chemistry, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

## GRAPHICAL ABSTRACT




**Abstract** Treatment of thiocarbonyl compounds with excess LDA leads to the corresponding thiols or sulfides depending on the work-up procedure. The mechanistic scenario for this unusual reduction pathway is discussed.

**Keywords** thioketones; LDA; hydride transfer; sulfides; reaction mechanisms

Address correspondence to Grzegorz Mlostoń, Department of Organic and Applied Chemistry, University of Łódź, Tamka 12, 91-403 Łódź, Poland. E-mail: gmloston@uni.lodz.pl


## INTRODUCTION

Thioketones belong to the class of reactive dipolarophiles (so-called superdipolarophiles) widely applied for the synthesis of numerous *S*-heterocyclic systems.<sup>1</sup> However, their behavior towards lithiated agents is by far less well known. As a part of our ongoing project focused on the exploration of thioketones in organic, materials, coordination, and biometalloorganic chemistry,<sup>2</sup> a series of model compounds of type **1** was recently shown to be suitable reaction partners in reactions with *C*-nucleophiles. For example, treatment of adamantanethione (**1a**) with lithiated methylphosphonate followed by methyl iodide or with methoxyallene anion provided the corresponding products, i.e., the phosphonylated sulfide **2**<sup>3</sup> and vinylthiirane derivative **3**,<sup>4</sup> respectively, as the result of exclusive *carbophilic* attack onto the C=S group (Scheme 1).



**Scheme 1** Reactions of adamantanethione (**1a**) with lithiated C-nucleophiles.


Within the studies on [3+2]-cycloadditions of thioketones with azomethine ylides, 1,3-thiazole-5(4*H*)-thione **4a** was selected as a model compound.<sup>5</sup> Unexpectedly, the treatment of a mixture of **4a** and trimethylamine oxide, used as the anticipated source of the parent azomethine ylide,<sup>6</sup> with LDA at 0 °C provided only the reduction product **5a**. Further experiments with **4a** confirmed the unusual potential of LDA for the reduction of the C=S group. Thus, depending on the work-up procedure, the corresponding thiol **5a** or methylsulfide **6a** were obtained in high yields<sup>7</sup> (Scheme 2).



**Scheme 2** Reduction of 1,3-thiazole-5(4*H*)-thione **4a** with LDA.


## RESULTS AND DISCUSSION

Following the protocol established for compound **4a**, a spirocyclic 1,3-thiazole-5(4*H*)-thione of type **4** and a series of thioketones **1** were smoothly reduced with excess of LDA to give the respective thiolates, which after trapping with methyl iodide as the electrophile provided the expected products **6b** and **7a-f**, respectively, in high yields (Figure 1). Analytically pure samples of products were obtained after chromatographic purification in 40–80% yields. Hence, as shown in Figure 1, dithioacetals **6a-b**, as well as cycloaliphatic (**7a-b**) and aromatic (**7c-f**) sulfides including unique diferrocenyl (**7e**) and hetaryl (**7f**) derivatives are available by the presented method, though the yields in the latter cases are rather low.<sup>7</sup>



**Figure 1** Products **6** and **7** prepared by the reduction of the respective thiocarbonyl substrates.

Although lithium amides are known in the first line as a strong bases, often used for the deprotonation of CH-acidic compounds, reducing properties of LDA and its analogs have also been reported.<sup>8</sup> For example, LDA-induced conversion of nitroarenes to the corresponding aromatic amines and azoxyarenes via a single electron transfer (SET) mechanism was described.<sup>8a</sup> **An analogous reaction pathway was postulated for the observed formation of sulfides from 2,2-diaryl-1,3-dithiolanes via the in situ generated aromatic thioketones.<sup>8b</sup>** On the other hand, treatment of 4-fluorotoluene with LDA in diethyl ether gave, among other products, a mixture of *meta*- and *para*-ethyl(2-tolylethyl)amine.<sup>9a</sup> A reaction mechanism via hydride transfer from LiNEt<sub>2</sub> to the intermediate aryne and subsequent addition of the aryl anion to the formed imine was proposed. Furthermore, the reduction of benzophenone with LiNEt<sub>2</sub> was interpreted as a hydride-transfer reaction via a six-membered transition state.<sup>9b</sup> A strong evidence for this reaction mechanism are enantioselective reductions of ketones with enantiomerically pure lithium alkyl phenyl amides<sup>9c</sup> and lithium dialkylamides.<sup>9d</sup> This pathway resembles that of reactions of Grignard reagents with sterically demanding carbonyl compounds as well as Meerwein-Schmidt-Ponndorf-Verley reductions of ketones.<sup>10</sup>



**Scheme 3** Postulated transition state (**A**) and hydride shift leading to thiolate **B**.

Based on the reports discussed above we suggest, that LDA in the reaction with non-enolizable thioketones acts as a hydride donor. Hydride transfer via the 6-membered

transition state **A** leads then to thiolate **B**, which can be protonated or trapped with appropriate electrophiles. The respective imine **C** is formed as a side-product (Scheme 3).<sup>[1]</sup> However, the competitive SET mechanism, especially in the case of hetaryl substituted thioketone **1f**, can't be ruled out.

## ACKNOWLEDGEMENTS

Authors acknowledge financial support by the National Science Center (PL-Cracow) Grant Maestro-3 no. 2012/06/A/ST5/00219. M.J. also thanks the University of Łódź Foundation. Stimulating discussion with Professor D. Seebach is also acknowledged.

## REFERENCES

- [1] (a) Fišera, L.; Huisgen, R.; Kalwinski, I.; Langhals, E.; Li, X.; Mlostoń, G.; Polborn, K.; Rapp, J.; Sicking, W.; Sustmann, R. *Pure Appl. Chem.* **1996**, 68, 789–798; (b) Mlostoń, G.; Heimgartner, H. In: *Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products* (Eds.: Padwa, A.; Pearson, W. H.), Wiley, New York, 2002, pp. 315–360; (c) Mlostoń, G.; Heimgartner H. In: *Targets in Heterocyclic Systems*, Vol. 10 (Eds.: Attanasi, O. A.; Spinelli, D.), Italian Society of Chemistry, 2006, pp. 266–300.
- [2] (a) Mlostoń, G.; Urbaniak, K.; Gębicki, K.; Grzelak, P.; Heimgartner, H. *Heteroatom Chem.*, **2014**, 25, 548–555; (b) Mlostoń, G.; Urbaniak, K.; Szychowska, A.; Linden, A.; Heimgartner, H. *Heterocycles*, in press (DOI: 10.3987/COM-14-S(K)56); (c) Mlostoń, G.; Pipiak, P.; Heimgartner H. *Helv. Chim. Acta*, in press (DOI: 10.1002/hlca.201400334)
- [3] Mlostoń, G.; Obijalska, E.; Urbaniak, K.; Heimgartner, H. *Heteroatom Chem.* **2008**, 19, 182–187.
- [4] Jasiński, M.; Mlostoń, G.; Stolarski, M.; Costa, W.; Domínguez, M.; Reissig, H.-U. *Chem. Asian J.* **2014**, 9, 2641–2648.
- [5] (a) Gebert, A.; Linden, A.; Mlostoń, G.; Heimgartner, H. *Heterocycles* **2002**, 56, 393–402; (b) Gebert, A.; Mlostoń, G.; Linden, A.; Heimgartner, H. *Pol. J. Chem.* **2003**, 77, 157–167; (c) Gebert, A.; Linden, A.; Mlostoń, G.; Heimgartner, H. *Pol. J. Chem.* **2003**, 77, 867–879.
- [6] (a) Beugelmans, R.; Negron, G.; Roussi, G. *J. Chem. Soc., Chem. Commun.* **1983**, 31–32; (b) Beugelmans, R.; Chastanet, J.; Roussi, G. *Heterocycles* **1987**, 26, 3197–3202; (c) Chastanet, J.; Roussi, G. *J. Org. Chem.* **1988**, 53, 3808–3812.
- [7] Gebert, A.; Jasiński, M.; Mlostoń, G.; Heimgartner, H. *Helv. Chim. Acta* **2014**, 97, 931–938.
- [8] (a) Wang, A.; Biehl, E. *Arkivoc* **2002**, (i), 71–75; (b) Ikehira, H.; Tanimoto, S.; Oida, T.; Okano, M. *J. Org. Chem.* **1983**, 48, 1120–1122; (c) Ashby, E. C.; Deshpande, A. K.; Patil, G. S. *J. Org. Chem.* **1995**, 60, 663–672; (d) De Kimpe, N.; Yao, Z. P.; Schamp, N. *Tetrahedron Lett.* **1986**, 27, 1707–1710; (e) Newcomb, M.; Burchill, M. T. *J. Am. Chem. Soc.* **1984**, 106, 8276–8282; (f) Newkome, G. R.; Hager, D. C. *J. Org. Chem.* **1982**, 47, 599–601; (g) Shen, C. C.; Ainsworth, C. *Tetrahedron Lett.* **1979**, 20, 89–92. For a review see: Majewski, M.; Cleave, D. M. *J. Organomet. Chem.* **1994**, 470, 1–16.
- [9] (a) Wittig, G.; Rentzea, C. N.; Rentzea, M. *Liebigs Ann. Chem.* **1971**, 744, 8–14; (b) Wittig, G.; Schmidt, H.-J.; Renner, H. *Chem. Ber.* **1962**, 95, 2377–2383; (c) Wittig,

G.; Thiele, U. *Liebigs Ann. Chem.* **1969**, 726, 1–12; (d) Takeda, K.; Ohnishi, Y.; Koizumi, T. *Org. Lett.* **1999**, 1, 237–239.

[10] (a) Maruyama, K.; Katagiri, T. *J. Phys. Org. Chem.* **1989**, 2, 205–213; (b) Campbell, E. J.; Zhou, H.; Nguyen, S.-B. T. *Angew. Chem., Int. Ed.* **2002**, 41, 1020–1022; *Angew. Chem.* **2002**, 114, 1062–1064.

[11] A small amount of the imine **C** was identified in the  $^1\text{H}$  NMR spectra of crude mixtures obtained with both adamantanthione (**1a**) and thiobenzophenone (**1c**); a characteristic septet attributed to the  $\text{CHMe}_2$  group was found at 3.77 ppm.<sup>12</sup> In addition, no incorporation of a deuterium atom was observed, when the reaction of LDA with **4a** was performed in  $\text{THF-}d_8$ .<sup>7</sup>

[12] Bunnelle, W. H.; Singam, P. R.; Narayanan, B. A.; Bradshaw, C. W.; Liou, J. S. *Synthesis* **1997**, 439–442.